

Agricultural Marketing Service, USDA

§ 29.3040

§ 29.3030 Greenish (V).

A color term applied to greenish-tinged tobacco. Any leaf which has a greenish tinge or a pale green color affecting 20 percent or more of its surface may be described as greenish. (See Rule 17.)

[24 FR 8771, Oct. 29, 1959. Redesignated at 49 FR 16757, Apr. 20, 1984]

§ 29.3031 Group.

A division of a type covering closely related grades based on certain characteristics which are related to stalk position or the general quality of the tobacco. Groups in Burley, Types 31 and 93, are as follows: Flyings (X), Lugs or Cutters (C), Leaf (B), Tips (T), Mixed (M), Nondescript (N), and Scrap (S).

[24 FR 8771, Oct. 29, 1959. Redesignated and amended at 49 FR 16757, Apr. 20, 1984]

§ 29.3032 Injury.

Hurt or impairment from any cause except the fungous or bacterial diseases which attack tobacco in its cured state. (See definition of Damage.) Injury to tobacco may be caused by field diseases, insects, or weather conditions; insecticides, fungicides, or cell growth inhibitors; nutritional deficiencies or excesses; or improper fertilizing, harvesting, curing, or handling. Injured tobacco includes dead, burnt, hail-cut, torn, broken, frost-bitten, sunburned, sunscalded, scorched, fire-killed, bulk-burnt, steam-burnt, barn-burnt, house-burnt, bleached, bruised, discolored, or deformed leaves; or tobacco affected by wildfire, rust, frog-eye, mosaic, root rot, wilt, black shank, or other diseases. (See Elements of quality and Rule 14.)

[24 FR 8771, Oct. 29, 1959. Redesignated at 49 FR 16757, Apr. 20, 1984]

§ 29.3033 Leaf.

Whole, unstemmed leaf. Leaf, when applied to tobacco in strip form, shall describe the divided unit of a whole leaf.

[49 FR 16758, Apr. 20, 1984]

§ 29.3034 Leaf scrap.

A by-product of unstemmed tobacco. Leaf scrap results from handling

unstemmed tobacco and consists of loose and tangled whole or broken leaves.

[24 FR 8771, Oct. 29, 1959. Redesignated at 49 FR 16758, Apr. 20, 1984]

§ 29.3035 Leaf structure.

The cell development of a leaf as indicated by its porosity or solidity. (See Elements of quality.)

[24 FR 8771, Oct. 29, 1959. Redesignated at 49 FR 16758, Apr. 20, 1984]

§ 29.3036 Leaf surface.

The smoothness or roughness of the web or lamina of a tobacco leaf. Leaf surface is affected to some extent by the size and shrinkage of the veins or fibers. (See Elements of quality.)

[24 FR 8771, Oct. 29, 1959. Redesignated at 49 FR 16758, Apr. 20, 1984]

§ 29.3037 Length.

The linear measurement of cured tobacco leaves from the butt of the midrib to the extreme tip. Length, as an element of quality, does not apply to tobacco in strip form. (See Elements of quality.)

[24 FR 8771, Oct. 29, 1959. Redesignated and amended at 49 FR 16757, 16758, Apr. 20, 1984]

§ 29.3038 Lot.

A pile, basket, bulk, bale or bales, sheet, case hogshead, tierce, package, or other definite package unit.

[48 FR 40366, Sept. 7, 1983. Redesignated at 49 FR 16758, Apr. 20, 1984]

§ 29.3039 Maturity.

The degree of ripeness. Tobacco is mature when it reaches its prime state of development. The extremes are expressed as immature and mellow. (See Elements of quality.)

[24 FR 8771, Oct. 29, 1959; 24 FR 9121, Nov. 10, 1959. Redesignated at 49 FR 16758, Apr. 20, 1984]

§ 29.3040 Mixed color (M).

Distinctly different colors of the type mingled together. (See Rule 16.)

[24 FR 8771, Oct. 29, 1959. Redesignated at 49 FR 16758, Apr. 20, 1984]